

# Cambridge International AS & A Level

CHEMISTRY 9701/13

Paper 1 Multiple Choice May/June 2020

1 hour

You must answer on the multiple choice answer sheet.

You will need: Multiple choice answer sheet

Soft clean eraser

Soft pencil (type B or HB is recommended)

Data booklet

#### INSTRUCTIONS

There are forty questions on this paper. Answer all questions.

- For each question there are four possible answers A, B, C and D. Choose the one you consider correct
  and record your choice in soft pencil on the multiple choice answer sheet.
- Follow the instructions on the multiple choice answer sheet.
- Write in soft pencil.
- Write your name, centre number and candidate number on the multiple choice answer sheet in the spaces provided unless this has been done for you.
- Do not use correction fluid.
- Do not write on any bar codes.
- You may use a calculator.

#### INFORMATION

- The total mark for this paper is 40.
- Each correct answer will score one mark. A mark will not be deducted for a wrong answer.
- Any rough working should be done on this question paper.

#### I have used following colour codes:

- Blue colour used for facts or reasoning
- · Red colour used for steps part of calculations along with its explanation
- Black colour used for right answer choice

This document has 16 pages. Blank pages are indicated.

IB20 06\_9701\_13/3RP © UCLES 2020

[Turn over

#### Section A

For each question there are four possible answers A, B, C and D. Choose the one you consider to be


Use of the Data Booklet may be appropriate for some questions.

Which particle has equal numbers of protons and neutrons and an electronic structure of  $1s^22s^22p^63s^23p^6$ ?





Which molecule contains six bonding electrons?



Solid carbon dioxide, CO<sub>2</sub>, is similar to solid iodine, I<sub>2</sub>, in its structure.

Which statement about solid CO<sub>2</sub> and solid SiO<sub>2</sub> is correct?

- Both solid CO<sub>2</sub> and solid SiO<sub>2</sub> exist in a lattice structure.
- Both solid CO2 and solid SiO2 have a simple molecular structure. SiO2 has giant molecular structure
- Both solid  $CO_2$  and solid  $SiO_2$  have atoms joined by single covalent bonds.
- Both solid CO<sub>2</sub> and solid SiO<sub>2</sub> change spontaneously to gas at s.t.p..
- The enthalpy changes of two reactions are shown.

$$K_2CO_3(s) + 2HCl(aq) \rightarrow 2KCl(aq) + H_2O(l) + CO_2(g)$$
  $\Delta H = -34.0 \text{ kJ mol}^{-1}$ 

$$KHCO_3(s) + HCl(aq) \rightarrow KCl(aq) + H_2O(l) + CO_2(g)$$
  $\Delta H = +32.8 \text{ kJ mol}^{-1}$ 

What is the enthalpy change for the reaction shown?

$$2KHCO_3(s) \rightarrow K_2CO_3(s) + H_2O(l) + CO_2(l)$$

$$2KHCO_{3}(s) \rightarrow K_{2}CO_{3}(s) + H_{2}O(1) + CO_{2}(g)$$

$$2 + 1Cl + 2KHCO_{3} \xrightarrow{2} K_{2}CO_{3} + H_{2}O + CO_{2} + 2HCl$$

$$2(+32.8) \times K_{2}CO_{3} + H_{2}O + CO_{2} + 2HCl$$

$$2(+32.8) \times K_{2}CO_{3} + H_{2}O + CO_{2} + 2HCl$$

В

-31.6 kJ mol<sup>-1</sup> 1.2 kJ mol<sup>-1</sup>

© UCLES 2020

Nitrogen reacts with oxygen to form nitrogen monoxide, NO, and nitrogen dioxide, NO<sub>2</sub>. Nitrogen dioxide reacts with water and with hydroxide ions.

What can be deduced using only the information from these equations?

HNO<sub>2</sub> is a strong acid.

HNO₃ is a weak acid.

NO<sub>2</sub> is a neutral gas.

NO is a reducing agent. Oxidises itself from +2 to +4 & reduces 02 from 0 to -2

Which solution has the lowest pH value?

A 0.01 mol dm<sup>-3</sup> butanoic acid

0.01 mol dm<sup>-3</sup> butanoic acid  $C_4H_4\omega_0H\rightarrow [C_4H_4\omega_0][H^+]\rightarrow [H^+]=0.01 \, \text{mol} \, / \text{dm}^3 \left(pH=2\right)$ 0.01 mol dm<sup>-3</sup> ethanoic acid  $CH_3\omega_0H\rightarrow [CH_3\omega_0][H^+]\rightarrow [H^+]=0.01 \, \text{mol} \, / \text{dm}^3 \left(pH=2\right)$ 0.01 mol dm<sup>-3</sup> hydrochloric acid  $HCL\rightarrow [H^+][CL]\rightarrow [H^+]=0.01 \, \text{mol} \, / \text{dm}^3 \left(pH=2\right)$   $U_2SO_4\rightarrow 2[H]^+[SO_4^{2-1}]\rightarrow [H^+]=2\times 0.01 \, \text{mol} \, / \text{dm}^3 =0.02 \, \text{mol} \, / \text{dm}^3 \left(pH=1.7\right)$ 0.01 mol dm<sup>-3</sup> sulfuric acid

The element sulfur produces a mass spectrum with the following peaks.

m/e value of peak	relative abundance
32	95.02
33	0.76
34	4.20
36	0.02

To calculate relative atomic man, multiply each relative abundance with its m/e value and take overall sum followed by division with 100.

Which relative atomic mass of sulfur can be calculated from these data, given to four significant figures?

32.07

32.08

32.09

32.10

[Turn over © UCLES 2020 9701/13/M/J/20



- 8 What is the electronic configuration of an isolated Ni<sup>2+</sup> ion?
  - $1s^22s^22p^63s^23p^63d^64s^2$
- "Ni → 15²25²2p°35°3p°45°3d"

B 1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>3s<sup>2</sup>3p<sup>6</sup>3d<sup>7</sup>4s<sup>1</sup>

C 1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>3s<sup>2</sup>3p<sup>6</sup>3d<sup>10</sup>4s<sup>2</sup>


D 1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>3s<sup>2</sup>3p<sup>6</sup>3d<sup>8</sup>

When electrons are being removed, trats done by first removing from highest energy level which is 4s in this case hance first 2 electrons will be removed from 4s

- At 200 °C aluminium chloride is a gas with  $M_r = 267$ .  $A_{12}Cl_6$

What is the number of covalent bonds, dative covalent bonds and lone pairs of electrons in one molecule of aluminium chloride at 200 °C?  $\bullet = A[ \chi = C]$ 

	covalent bonds	dative covalent bonds	lone pairs
Α	6 /	2 🗸	0
B	41°C16	AL*CL2	√16 *×
C	6	2 /	18
D	3	0	9



10 When solid KClO<sub>3</sub> is heated in the absence of air, a mixture of two chlorine compounds in the mole ratio of 3:1 is formed. Chlorine is the only element whose oxidation number changes in this reaction.

What could be the oxidation numbers of chlorine in the two compounds that are formed?

We die from the two compounds that are formed?

\*\*eduction\*\*

- +3 and -1
- +6 and +4 (c) +7 and -1
- +7 and +1

11 Two reactions are shown.

reaction 1 
$$X_2(g) + Y_2(g) \rightleftharpoons 2XY(g)$$

reaction 2 
$$XY(g) \rightleftharpoons \frac{1}{2}X_2(g) + \frac{1}{2}Y_2(g)$$

The equilibrium constant,  $K_p$ , for reaction 1 is 0.0052.

What is  $K_p$  for reaction 2?

A 
$$2.6 \times 10^{-3}$$
 B  $13.9$  C  $192.3$  D  $384.6$ 

$$\frac{\left(XY\right)^{2}}{\left(X_{2}\right)\left[Y_{2}\right]} = 0.0052$$

$$\frac{\left(X_{2}\right)\left[Y_{2}\right]}{\left(XY\right)^{2}} = \frac{2500}{13}$$

$$\frac{\left(X_{2}\right)\left[Y_{2}\right]}{\left(XY\right)^{2}} = \frac{2500}{13}$$

$$\frac{\left(X_{2}\right)\left[Y_{2}\right]}{\left(XY\right)^{2}} = \frac{2500}{13}$$

While Cl reduces from

Take reciprocal of this lets root it

12 Compound T is a white cr	rvstalline	solia
-----------------------------	------------	-------


When a sample of compound T is mixed with aqueous sodium hydroxide and heated, a gas is produced which turns damp red litmus paper blue.

Further testing of a solution of compound T with aqueous barium chloride produces a dense white precipitate which does not dissolve when dilute hydrochloric acid is added to the mixture.

	Wh	at is the identity of compound T?
	Α	at is the identity of compound 1?  + NaDH -> NH3 validates 1 statement but (NH4)203+ Bacl2  ammonium carbonate gives Bacl3 which reacts with dilute HCl
	B	
	×	sodium carbonate + NaDH -> NH, validates 1st statement but (NH) SD.
	Ø	sodium carbonate + NaDH -> NH, validales 1st statement but (NH,) SD, when reacts with Bacl, gives Baso, that is not soluble in dilule HCI
13	C a Wh	ich property explains the trend in volatility of the elements going down Group 17?
	Α	decreasing covalent bond strength
	В	decreasing van der Waals' forces Volatility decreases down Group / due to Jincrease in number of electrons
	С	increasing covalent bond strength which lead to increased vander Waal forces
(		increasing van der Waals' forces
14		e statements apply to the elements in Group 2.
	vvn	ich statement is correct?
(		As atomic number increases, ionic radius increases.  As atomic number increases, reducing ability decreases.
	В	As atomic number increases, reducing ability decreases. X
	С	As atomic number increases, first ionisation energy increases. X down the group, i.e decreases
	D	As atomic number increases, first ionisation energy increases. X down the group, i.e decreases  As atomic radius increases, first ionisation energy increases. Xie decreases as electron gets farther from nucleus
15	Wh	ich element, when burned in oxygen, can form an oxide that is a reducing agent? Hat oxidises itself
	Α	Na B Mg C Al (D) S+4 +6
	t	$Na_10$ Mg0 $AY_1O_3$ $SO_2/SO_3$
		l 🔥


16	Nitrog	en oxides are removed of a catalyst in a catalyti		es of internal combustion engines b	y the
		row is correct?	C0+ NO	reduction in presence of platinum catalyst	
		change in oxidation number of nitrogen	type of catalyst	of platinum catalyst	
	(A)	decrease ✓	heterogeneous V		
	В	decrease 🗸	homogeneous		
	c	increase	heterogeneous		
	D	increase	homogeneous		
17	which	dissolves in an excess o	r nitrate to aqueous bar f dilute aqueous ammoni	$\rightarrow$ AgCl + Ba(N0 <sub>3</sub> ) <sub>2</sub> Actinum chloride produces a white precipitate to form a colourless solution.  AgCl + NH <sub>3</sub> $\longrightarrow$ Cl + Ag(NH <sub>3</sub> )  the colourless solution produces a	pitate
	What			HNO	3
(			Ba(NO <sub>3</sub> ) <sub>2</sub>	D NH4NO3 Agi	îl(s)
18	Which	property shows an <b>incre</b>	ease from calcium to bar	rium going down Group 2?	
(		ne ease of decomposition ne solubility of the hydrox		operative 1 down the group	
		ne solubility of the sulfate		(DIL)	
		ne volume of hydrogen gi	U	1	
19				form an alkaline solution. Solution	0 are alboline
/		statement about the <b>chl</b>		prrect? Nacl or MgCl <sub>2</sub>	
(		conducts electricity when			
		has a melting point of les	ss than 100°C.	CH3. B	r 2 43
		has covalent bonding.		$C=C$ + $+Br \rightarrow -C -C$	1-c-c-c
Function 20	ral/chair	reacts rapidly with cold v Aposition cis/b ural and stereoisomerism	rans Epollical	when answering this question. $C_2H_5$	or - C - C - C
	When	trans-pent-2-ene reacts	with HBr, how many diffe	ferent products can form?	Br
	<b>A</b> 1	<b>B</b> 2	<b>C</b> 3	<b>D</b> 4	
© U(	CLES 202	0	9701/13/M/J/20		

21 Ester P has the following structural formula.



Which compounds are produced when P is hydrolysed using dilute hydrochloric acid?

- A CH<sub>3</sub>COC1 and (CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>CH<sub>2</sub>OH
- B CH<sub>3</sub>CH<sub>2</sub>OH and (CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>CO<sub>2</sub>H
- CH3CO2H and (CH3)2CHCH2CO2H



22 There are many non-cyclic alcohols that cannot be oxidised by warm acidified  $MnO_4^-$  ions. Alcohol X is the member of this set of alcohols with the **lowest** molecular mass.

How many moles of oxygen are required for the complete combustion of 1.0 mol of alcohol X?

- A 3.5 mol
- **B** 4.5 mol
- D 6.5 mol  $C_4 \parallel_{10} 0$  + $GD_2 \longrightarrow 4CO_2 +5 \parallel_2 0$


23 Butanoic acid can be produced from 1-bromopropane in two steps using reagents V and W as maly VMD can add, a

		on ex	tra carbon	
3 carbons	reagent V		reagent W	4 carbons
1-bromopropane		compound Q	$\longrightarrow$	butanoic acid


What could be reagents V and W?

	V	W
A	KCN in ethanol	HCl(aq) 🗸
В	KCN in ethanol 🗸	NaOH(aq)
С	NH₃ in ethanol	HCl(aq)
D	NaOH(aq)	H <sup>+</sup> /Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup> (aq)

© UCLES 2020 9701/13/M/J/20 [Turn over



- T and U are stereoisomers.
- **B** T can be distinguished from U by the use of alkaline aqueous iodine.
- T can be reduced by LiAIH4 but not by NaBH4. both are veducing agouls
- U can be formed by the oxidation of 3-methylbutan-1-ol.
- 25 The diagram shows the infrared spectrum of an organic compound.



What could be the identity of this compound?

propan-1-of no strong peak btw 3200 & 3650 propanal no strong peak at 1670-1740

c propanoic acid propanone no strong peak at 1670-1740

26 Which reagent reacts with both of the -OH groups in lactic acid, CH<sub>3</sub>CH(OH)CO<sub>2</sub>H?

sodium hydroxide only acid

© UCLES 2020

9701/13/M/J/20

27 1,2-dibromopropane can be made from 1-bromopropane in two steps.

Which row is correct?

	step 1	step 2
Α	addition	substitution
В	elimination	addition
X	hydrolysis	elimination
X	substitution	hydrolysis

28 2-methylbut-2-ene reacts with HBr(g) to form two isomeric products. During the reaction two Which diagram shows the more stable of the two positively charged intermediates?  $\frac{cH_3}{c} = \frac{cH_3}{c} =$ 





29 The ester ethyl methanoate is prepared in a school laboratory by reacting a carboxylic acid with an alcohol.

During the reaction, only 50.0% of the alcohol is converted into the ester.

Which mass of alcohol is needed to prepare 10.0 g of the ester?



A 3.11g B 8.65g C 12.4g D 32.2g that is converted H<sub>2</sub>Ot  $C_3H_6O_2 \longrightarrow CH_2O_2 + C_2H_6O$  mass of alwhol that is converted but various  $O \cdot 13C_2 = \frac{10}{70} \cdot 13S_2 = \frac{10}{70} \cdot 13S_2$ 

## 30 Compound X has the structure shown.

Which type of carbonyl group is present and how many chiral centres are there in one molecule of X?

	carbonyl group	chiral centres
Α	aldehyde	0
В	aldehyde	1
С	ketone 🗸	0
	ketone 🗸	1

#### Section B

For each of the questions in this section, one or more of the three numbered statements 1 to 3 may be correct.

Decide whether each of the statements is or is not correct (you may find it helpful to put a tick against the statements that you consider to be correct).

The responses A to D should be selected on the basis of

	Α	В	С	D
1	, <b>2</b> and <b>3</b> are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

Use of the Data Booklet may be appropriate for some questions.

31 Which contain one mole of the underlined substance under room conditions?

I a balloon containing 24.0 dm3 of helium wolar volume of each mole of gas

✓ a block of <u>calcium carbonate</u> weighing 100.1 g Mr = | DD • |

4000 x 0.250 = 1 mg 4000 cm<sup>3</sup> of a 0.250 mol dm<sup>-3</sup> solution of sulfuric acid

32 Buckminsterfullerene is a fullerene allotrope of carbon.

Buckminsterfullerene is a giant covalent molecule. Buckminsterfullerene has delocalised electrons. like graphile

Buckminsterfullerene has strong intramolecular bonds.

33 Gaseous sodium ions can be formed from sodium atoms.

from sodium atoms. Atomisation ionisation Na(s) o Na(g) o Na(g) o Na(g)

Which quantities are required to calculate the enthalpy change of formation of Na<sup>+</sup>(g)?

first ionisation energy of sodium

enthalpy change of atomisation of sodium

enthalpy change of formation of sodium

[Turn over © UCLES 2020 9701/13/M/J/20

The responses A to D should be selected on the basis of

Α	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

34 The Haber process is used in industry to form ammonia from hydrogen and nitrogen.

$$3H_2 + N_2 \rightleftharpoons 2NH_3$$

Which statements about the activation energy for this process are correct?

The activation energy for the forward reaction is the same as the activation energy for the reverse reaction.

The activation energy for the reverse reaction is decreased by the addition of iron.

The activation energy is the minimum energy that colliding particles must possess in order to react.

activation energy for backward reaction

35 Strontium nitrate is heated strongly for several minutes.

Which statements are correct? A

A brown gas is produced. NO2

 $\nearrow$  A gas is produced that relights a glowing splint.  $\bigcirc_2$ 

→ A white powder remains after heating. Sro

36 When added to water, which oxides will **not** cause a change in pH?

$$4Al_2O_3$$
  $Al_2O_3+H_2O \longrightarrow insoluble$ 

© UCLES 2020

37 Propanal reacts with hydrogen cyanide to form 2-hydroxybutanenitrile. A suitable catalyst for this reaction is sodium cyanide.


Which statements about this catalysed reaction of propanal with hydrogen cyanide are correct?  ${f D}$ 



The sodium cyanide provides a stronger nucleophile than HCN.

The reaction can be classified as nucleophilic substitution. nucleophilic addition

The hydrogen cyanide molecule attacks the propanal molecule to form an intermediate ion.



38 A reaction mechanism is shown.

$$H_2C$$
 $CH_2$ 
 $H_2C$ 
 $CH_2$ 
 $CH_2$ 

Which statements about this reaction are correct? A

It is a substitution reaction.

OH⁻ behaves as a nucleophile.

Heterolytic bond fission is involved. C-CL & X-OH

39 On complete combustion, a sample of X produces 44 g of carbon dioxide and 27 g of water. On complete combustion, a sample of Y produces 44 g of carbon dioxide and 18 g of water. On complete combustion, a sample of Z produces 22g of carbon dioxide and 9g of water.

Which substances could be straight chain alkanes?



Straight chain alkanes are more stable thus give out more heat on combastion

Y and Z relatively produce less amount of heat

© UCLES 2020 [Turn over 9701/13/M/J/20

The responses A to D should be selected on the basis of

A	В	С	D
1, 2 and 3 are correct	1 and 2 only are correct	2 and 3 only are correct	1 only is correct

No other combination of statements is used as a correct response.

40 Which pairs are structural isomers of each other?

CH3CH2CH2CH2CO2H and CH3CH2CH2CO2CH2CH3 (functional isomer)

 $\textbf{3} \quad \mathsf{CH_3CH_2CH_0CH_0CH_3} \ \ \textbf{and} \ \ \mathsf{CH_3CH_2CH(OH)CH_2CH_3}$ 

## **BLANK PAGE**

### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.