



## Cambridge International AS & A Level

**CHEMISTRY**

**9701/12**


Paper 1 Multiple Choice

**February/March 2022**

**1 hour 15 minutes**

You must answer on the multiple choice answer sheet.

You will need: Multiple choice answer sheet  
Soft clean eraser  
Soft pencil (type B or HB is recommended)



### INSTRUCTIONS

- There are **forty** questions on this paper. Answer **all** questions.
- For each question there are four possible answers **A**, **B**, **C** and **D**. Choose the **one** you consider correct and record your choice in soft pencil on the multiple choice answer sheet.
- Follow the instructions on the multiple choice answer sheet.
- Write in soft pencil.
- Write your name, centre number and candidate number on the multiple choice answer sheet in the spaces provided unless this has been done for you.
- Do **not** use correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.

### INFORMATION

- The total mark for this paper is 40.
- Each correct answer will score one mark.
- Any rough working should be done on this question paper.
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

---

This document has **20** pages. Any blank pages are indicated.



1 The first ionisation energy of potassium, K, is  $418 \text{ kJ mol}^{-1}$ . The first ionisation energy of strontium, Sr, is  $548 \text{ kJ mol}^{-1}$ . *Since potassium & strontium are from the same period, their no. of shells remain the same except for the no. of protons which increase along the period.*

Which statement helps to explain why Sr has a greater first ionisation energy than K?

- The more the no. of protons, the greater the charge and hence increased ionis. energy*
- A The charge on a Sr nucleus is greater than the charge on a K nucleus.
- B The outer electron in a Sr atom experiences greater shielding than the outer electron in a K atom. *When shielding increases, ionisation energy must decrease*
- C The outer electron in a Sr atom experiences spin-pair repulsion. *in this case, i.e must be less*
- D The outer electron in a Sr atom is further from the nucleus than the outer electron in a K atom. *the greater the bond length, the lower the ionis. energy*

2 What is the electronic configuration of  $\text{Mg}^{2+}$ ?

- A  $1s^2 2s^2 2p^6$
- B  $1s^2 2s^2 2p^6 3s^2$
- C  $1s^2 2s^2 2p^6 3s^2 3p^2$
- D  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^2 4s^2$



*electrons are removed from shells that have higher energy.*

3 Compound X contains the elements C, H and O only.

2.00 g of X produces 4.00 g of carbon dioxide and 1.63 g of water when completely combusted.

What is the empirical formula of X?

- A  $\text{CHO}_2$  *0.04 moles would produce 0.04 moles of  $\text{CO}_2$*
- B  $\text{C}_2\text{H}_2\text{O}$  *0.048 moles would produce 0.075 moles of  $\text{CO}_2$*
- C  $\text{C}_2\text{H}_4\text{O}$  *0.045 moles would produce 0.091 moles of  $\text{CO}_2$*
- D  $\text{CH}_2\text{O}_2$

4 For which molecule is the dipole moment zero?



- A  $\text{CH}_3\text{Cl}$
- B  $\text{CH}_2\text{Cl}_2$
- C  $\text{CHCl}_3$
- D  $\text{CCl}_4$

*0.091 moles  $\text{CO}_2$*       *0.07056 moles  $\text{H}_2\text{O}$*

$$\text{C}_2\text{H}_4\text{O} + \frac{5}{2}\text{O}_2 \rightarrow 2\text{CO}_2 + 2\text{H}_2\text{O}$$

<sup>3</sup>  
*rule of dative bonding*



5 Which dot-and-cross diagram is correct for  $Al_2Cl_6$ ?



key

× = electrons from Al

○ = electrons from Cl



6 The boiling points of some hydrogen halides are shown.

hydrogen halide	boiling point / K
H-Cl	188
H-Br	206
H-I	238

*b.p. increasing*

What is the explanation for the trend in boiling point for the hydrogen halides from HCl to HI?

- The bond energies of the hydrogen halides increase from HCl to HI. *bond energy decreases from HCl to HI*
- B** There is an increase in the strength of the intermolecular forces of attraction from HCl to HI. *due to ↑ in no. of electrons*
- C The intermolecular hydrogen bonds become stronger from HCl to HI. *Hydrogen bonds form among O, N and F along with H*
- D There is an increase in the bond polarity from HCl to HI.

7 Elements X, Y and Z are all in the first two periods of the Periodic Table.

Their Pauling electronegativity values,  $E_N$ , are shown.

element	$E_N$
X	1.0
Y	2.1
Z	4.0

*the greater the  $E_N$  difference, the stronger the bond*

Substances exist with formulae XZ, YZ and Z<sub>2</sub>. *which has covalent bonding and intermolecular forces*

Which row puts these substances in order of increasing melting point?

	lowest melting point	→	highest melting point
<del>A</del>	XZ		Z <sub>2</sub> <del>X</del>
B	XZ		YZ
<b>C</b>	Z <sub>2</sub>		XZ ✓
D	Z <sub>2</sub>		YZ


8 The equation for reaction 1 is shown.



	$\Delta H_c^\ominus / \text{kJ mol}^{-1}$
$C_6H_{12}O_6$	a
$C_2H_5OH$	b

*when 1 mole of  $C_2H_5OH$  is completely combusted*

*enthalpy change of combustion*




What is the correct expression for the enthalpy change of reaction 1?

- A  $a + b$       B  $a - b$       C  $a + 2b$       **D**  $a - 2b$


$$a = x + 2b$$

$$x = a - 2b$$


- 9 Nitrogen monoxide is an atmospheric pollutant that is formed inside car engines by an endothermic reaction between nitrogen and oxygen.




Which diagram correctly represents the energy profile for this reaction?



- 10 Two half-equations are shown.



The equation for the reaction between manganate(VII) ions and sulfite ions is shown.





Which statements are correct?

- 1  $u = x = z$
- 2 Manganese is reduced to oxidation state +4.
- 3 Sulfur is oxidised from oxidation state +4 to +6.

- A** 1, 2 and 3    **B** 1 and 2 only    **C** 1 and 3 only    **D** 2 and 3 only

- 11 Hydrogen peroxide,  $\text{H}_2\text{O}_2$ , decomposes to form water and oxygen.


The reaction is catalysed by bromide ions.



Which row is correct?

	type of catalyst	in step 1
<del>A</del>	heterogeneous <del>X</del>	bromide ions are oxidised <del>✓</del>
<del>B</del>	heterogeneous	bromide ions are reduced <del>X</del>
<b>C</b>	homogeneous <i>same state as reactants</i>	bromide ions are oxidised <del>✓</del>
<del>D</del>	homogeneous	bromide ions are reduced <del>X</del>

- 12 Hydrogen and iodine react to form hydrogen iodide in an exothermic reaction. The equation is shown.



*equilibrium shifts to the side that has fewer no. of moles*



A  $1 \text{ m}^3$  reaction vessel contains  $\text{H}_2$ ,  $\text{I}_2$  and  $\text{HI}$  gases at equilibrium. The temperature is changed such that the total pressure in the  $1 \text{ m}^3$  vessel doubles.

*temperature is increased so backward reaction is favoured.*

What is the effect on the value of  $K_p$  and on the position of equilibrium?

	effect on the value of $K_p$	effect on the position of equilibrium
<b>A</b>	decreases	moves left
<del>B</del>	increases	moves right <del>X</del>
<del>C</del>	no change	moves left
<del>D</del>	no change	no change <del>X</del>

13 Diethylzinc,  $(\text{C}_2\text{H}_5)_2\text{Zn}$ , is added to  $\text{NaOH}(\text{aq})$ . Two reactions occur.



In these reactions, which compounds act as Brønsted–Lowry acids?


	reaction 1	reaction 2
A	$(\text{C}_2\text{H}_5)_2\text{Zn}$	$\text{H}_2\text{O}$
<b>B</b>	$\text{H}_2\text{O}$	$\text{H}_2\text{O}$
<del>C</del>	$\text{H}_2\text{O}$	$\text{ZnO}$
<del>D</del>	the reaction is not acid/base	$\text{ZnO}$

14 Which statement about atoms and molecules is correct?

- ~~A~~ The molecular formula of a compound is the simplest whole number ratio of atoms of each element in the compound. *that is empirical formula*
- B One mole of any substance contains  $6 \times 10^{23}$  atoms.
- ~~C~~ The relative atomic mass of an element is the ratio of the average mass of one atom of the element to the mass of an atom of carbon-12.
- D** The relative formula mass of a compound is the sum of the individual atomic masses of all the atoms in the formula.







15 The Boltzmann distribution for one mole of a gas at temperature  $T$  is shown.




One mole of the same gas is added, and the gas remains at temperature  $T$ . *concentration ↑* *rate of reaction ↑ but  $E_a$  remains the same*

Which dotted curve shows the distribution with the added gas?



- 16 In the reaction shown, the concentrations of both X and Y are reduced to half of their original values whilst keeping the total volume of the solution constant. *affects rate*



Simultaneously the temperature is increased from 298 K to 348 K. *affects rate*

Which prediction is definitely true?

- A A smaller proportion of collisions between particles of X and particles of Y will be successful. *only in the case of decreasing concentration*
- B The average kinetic energy of particles of X and particles of Y will increase.
- C The rate of the reaction will be unaffected.
- D The frequency of collisions between particles of X and particles of Y will halve. *only in the case of decreasing concentration*

- 17 A student investigated the chloride of a Period 3 element. This is what the student wrote down as a record.

*If dissolves in water then the pH should remain near 7.0*

The compound was a white crystalline solid. It dissolved easily in water to give a solution of pH 12. When placed in a test-tube and heated in a roaring Bunsen flame, the compound melted after several minutes heating.

*Na, Mg or Al bcz non metallic chlorides hydrolyse in water & form HCl*

What can be deduced from this record?

- A At least one of the recorded observations is incorrect.
- B The compound was magnesium chloride,  $MgCl_2$ . *white solid, dissolves in water*
- C The compound was phosphorus pentachloride,  $PCl_5$ . *does not dissolve in water, yellowish crystals*
- D The compound was sodium chloride,  $NaCl$ . *yellow*

- 18 The elements in Period 3 and their compounds show trends across the period from sodium to chlorine.

Which row is correct?

*metallic oxides → basic in nature*  
*non metallic oxides → acidic in nature*

	electronegativity of the elements	acid / base behaviour of the oxides of the elements
<input checked="" type="radio"/>	decreases <input checked="" type="checkbox"/>	basic → amphoteric → acidic
<input checked="" type="radio"/>	decreases <input checked="" type="checkbox"/>	acidic → amphoteric → basic <input checked="" type="checkbox"/>
<input checked="" type="radio"/> C	increases	basic → amphoteric → acidic <input checked="" type="checkbox"/>
D	increases	acidic → amphoteric → basic <input checked="" type="checkbox"/>

- 19 The table shows the melting points of  $\text{SiO}_2$  and  $\text{P}_4\text{O}_6$ .

oxide	$\text{SiO}_2$	$\text{P}_4\text{O}_6$
melting point/K	1883	297

Which statement explains the difference between the melting points of  $\text{SiO}_2$  and  $\text{P}_4\text{O}_6$ ?

- A The bonding of the oxides changes from ionic to covalent. *both have covalent*
- B The metallic character of the elements decreases across Period 3.
- C The oxidation number of the element increases from Si to P.
- D The structure changes from giant molecular to simple molecular.

- 20 Equal masses of  $\text{CaCO}_3$ ,  $\text{Ca}(\text{NO}_3)_2$ ,  $\text{BaCO}_3$  and  $\text{Ba}(\text{NO}_3)_2$  are thermally decomposed. The volume of gas produced in each experiment is measured under the same conditions.

Which compound will produce the greatest volume of gas?

- A  $\text{CaCO}_3$      B  $\text{Ca}(\text{NO}_3)_2$      C  $\text{BaCO}_3$      D  $\text{Ba}(\text{NO}_3)_2$

- 21 Which row gives correct comparisons between the solubilities of calcium hydroxide and barium hydroxide and the thermal stabilities of calcium carbonate and barium carbonate?

*solubility of Group II hydroxides increases down the group*

	solubility		thermal stability	
	calcium hydroxide	barium hydroxide	calcium carbonate	barium carbonate
<input checked="" type="checkbox"/> A	higher <input checked="" type="checkbox"/>	lower	higher	lower
<input checked="" type="checkbox"/> B	higher <input checked="" type="checkbox"/>	lower	lower	higher
<input type="checkbox"/> C	lower <input checked="" type="checkbox"/>	higher <input checked="" type="checkbox"/>	higher <input checked="" type="checkbox"/>	lower
<input checked="" type="checkbox"/> D	lower <input checked="" type="checkbox"/>	higher <input checked="" type="checkbox"/>	lower	higher

*stability ↑ down the group because more heat is required to decompose larger compounds*


- 22 Which statement relating to the elements in Group 17 and their compounds is correct?

- A Bromine will oxidise KI to form iodine. *More reactive halogen displaces less reactive halide ion from its compound*  $\text{Br}_2 + 2\text{KI} \rightarrow 2\text{KBr} + \text{I}_2$
- B Iodide ions react to form a white precipitate when added to silver nitrate solution. *yellow*
- C Bromide ions react to form a white precipitate when added to silver nitrate solution. *cream*
- D Chlorine reacts with hydrogen to form a colourless gas. *HCl*

- 23 An excess of chlorine was bubbled into 100 cm<sup>3</sup> of hot 6.0 mol dm<sup>-3</sup> sodium hydroxide.

How many moles of sodium chloride would be produced in the reaction?

- A 0.30     B 0.50    C 0.60    D 0.72




24 The product of the **Contact process** is Z.  $\text{SO}_2 + \text{O}_2 \rightarrow \text{SO}_3$

Which reaction or process leads to the formation of a gas that can **neutralise an aqueous solution of Z?**

- ~~A~~ atmospheric lightning gives NO  $\text{SO}_3 + \dots$   
 acidic basic  
~~B~~ combustion of fuel in an internal combustion engine  $\text{CO}_2$   
**C** the Haber process  $\text{NH}_3$  (basic) ✓  
~~D~~ thermal decomposition of Group 2 nitrates  $\text{NO}_2$

25 When ammonia,  $\text{NH}_3$ , is dissolved in water, a small concentration of ammonium ions,  $\text{NH}_4^+$ , is formed.

Which row is correct?



	number of electrons in one ammonium ion	change of the H-N-H angle from ammonia to the ammonium ion
<del>A</del>	8 <del>X</del>	decreases <del>X</del>
<del>B</del>	8 <del>X</del>	increases
<del>C</del>	10	decreases <del>X</del>
<b>D</b>	10	increases

these are only valence electrons  
 Nitrogen has 2 more electrons in inner most shell


26 In this question, **alkenes and cyclic alkanes** should be considered.

How many **structural isomers** of  $\text{C}_4\text{H}_8$  are there?

- A 3      B 4      **C** 5      D 6

27 Which compound will **decolourise**  $\text{Br}_2(\text{aq})$ ?

- ~~A~~  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$   
~~B~~  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CHO}$   
**C**  $\text{CH}_3\text{CHCHCH}_2\text{CH}_2\text{CH}_2\text{OH}$  is replaced by Br  
 D  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_3$




- 28 Alkenes react with aqueous hydrogen bromide. The reaction proceeds via an intermediate carbocation. The more stable the intermediate, the faster the reaction.

Which sequence correctly shows an increase in the speed of reaction of the alkenes with hydrogen bromide?

- (A) ethene, propene, 2-methylpropene  
 (B) 2-methylpropene, ethene, propene  
 (C) propene, ethene, 2-methylpropene  
 (D) propene, 2-methylpropene, ethene


addition reaction  
 stable intermediate forms which is only possible with secondary or tertiary structures with tertiary structure being most stable



- 29 A reaction occurs when a sample of 1-chloropropane is heated under reflux with sodium hydroxide dissolved in ethanol.

Which row is correct?


	type of reaction	name of product
A	elimination	propan-1-ol
(B)	elimination	propene
C	substitution	propan-1-ol
D	substitution	propene



If NaOH(aq) is used, C<sub>3</sub>H<sub>7</sub>OH forms where it's a substitution reaction

- 30 The diagram shows the structures of three halogenoalkanes.

Stable structures undergo S<sub>N</sub>2 mechanism which are tertiary structures



P, Q and R can all be hydrolysed. halides replaced by OH

Which row is correct?

	relative speed of hydrolysis		mechanism of hydrolysis	
	Q	R	P	Q
(A)	fast	slow	S <sub>N</sub> 1 ✓	S <sub>N</sub> 2
B	fast	slow	S <sub>N</sub> 2	S <sub>N</sub> 1 X
C	slow X	fast	S <sub>N</sub> 1	S <sub>N</sub> 2
D	slow X	fast	S <sub>N</sub> 2	S <sub>N</sub> 1

difference is in the bond energy which is greater of C-Cl the greater the bond energy, the slower is the reaction

13

*completely oxidised**oxidising agent*

- 31 A sample of <sup>0.05 moles</sup> 2.30 g of ethanol is mixed with an excess of aqueous acidified potassium dichromate(VI). The reaction mixture is boiled under reflux for one hour. The required organic product is then collected by distillation. The yield of product is 60.0%.  
*although distillation gives aldehyde, reflux will not help*  
 Which mass of product is collected?

A 1.32 g

B 1.38 g

 C 1.80 g



D 3.00 g



$$0.03 \times (12 + 3 + 12 + 32 + 1)$$

*0.03 moles of oxidised substance*

- 32 The structure of tartaric acid is shown.



Four moles of substance X react with one mole of tartaric acid.


What could be substance X?

- A sodium *reacts with both OH & COOH groups*  
 B sodium carbonate *reacts with only COOH & 2 moles of Na would react*  
 C sodium hydrogencarbonate  
 D sodium hydroxide *only 2 moles of Na react*

- 33 Which compound gives both:

- an orange precipitate with 2,4-DNPH reagent *both aldehydes & ketones do*
- and a yellow precipitate with alkaline  $I_2(aq)$ ? *only ketones, secondary alcohols or ethanal would*

- ethanol  
 methanal  
 propanal  
 D propanone



Which reactions would give the product shown?

- A both reactions
- B reaction 1 only
- C reaction 2 only
- D neither reaction

35 Compound X contains a <sup>-COO group</sup> single ester group.


X contains 27.6% by mass of oxygen.

Which pair of products could be produced by the hydrolysis of X?


- A butan-1-ol and ethanoic acid *butylethanoate*  $\text{CH}_3\text{COOC}_4\text{H}_9$  has  $\frac{32}{116} \times 100\% = 27.6\%$  of  $\text{O}_2$
- B ethanol and propanoic acid *ethyl propanoate*  $\text{C}_2\text{H}_5\text{COOC}_2\text{H}_5$  has  $\frac{32}{90} \times 100\% = 35.6\%$  of  $\text{O}_2$
- C methanol and methanoic acid *methyl methanoate*  $\text{HCOOCH}_3$  has  $\frac{32}{60} \times 100\% = 53.3\%$  of  $\text{O}_2$
- D propan-2-ol and butanoic acid *prop-2-yl butanoate*  $\text{C}_3\text{H}_7\text{COOC}(\text{CH}_3)_2$  has  $\frac{32}{130} \times 100\% = 24.6\%$  of  $\text{O}_2$

36 What is the least number of carbon atoms in a non-cyclic alkane molecule that has a chiral centre?

- A 7
- B 8
- C 9
- D 10



37 A molecule of a polymer contains the sequence shown.



Which monomer could produce this polymer by addition polymerisation?


- A  $\text{CHCl}=\text{CHCl}$
- B  $\text{CH}_2=\text{CHCl}$
- C  $\text{CH}_3\text{CCl}=\text{CHCl}$
- D  $\text{CH}_3\text{CCl}=\text{CH}_2$

- 38 Compound Y is heated with a mild oxidising agent. One of the products of the reaction reacts with hydrogen cyanide forming 2-hydroxybutanenitrile.

What is compound Y?

- HCN (addition)*
- A butan-1-ol  $\rightarrow$   $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH} \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CHO} \xrightarrow{\text{HCN}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{CN}$  hydroxybutane nitrile
- B butan-2-ol  $\rightarrow$   $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{C}(\text{O})\text{CH}_3 \xrightarrow{\text{HCN}} \text{CH}_3\text{CH}_2\text{C}(\text{OH})(\text{CN})\text{CH}_3$  2-hydroxybutane 2-nitrile
- C propan-1-ol  $\rightarrow$   $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} \rightarrow \text{CH}_3\text{CH}_2\text{CHO} \xrightarrow{\text{HCN}} \text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CN}$  2-hydroxy propano nitrile
- D propan-2-ol  $\rightarrow$   $\text{CH}_3\text{CH}(\text{OH})\text{CH}_3 \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_3 \xrightarrow{\text{HCN}} \text{CH}_3\text{C}(\text{OH})(\text{CN})\text{CH}_3$  2-hydroxy pentane 2-nitrile

- 39 The diagrams show the structures of lycopene and  $\beta$ -carotene.







When lycopene is converted into  $\beta$ -carotene, what is the gain or loss of hydrogen atoms per molecule?

- A 4 gained
- B 2 gained
- C no change
- D 2 lost



- 40 Which diagram shows the infrared spectrum of a compound that contains both a C=O and an O-H group? 2500-3650



bond	functional group containing the bond	characteristic infrared absorption range (in wavenumbers) / cm <sup>-1</sup>
C-O	hydroxy, ester	1040-1300
C=C	aromatic compound, alkene	1500-1680
C=O	amide carbonyl, carboxyl ester	1640-1690 1670-1740 1710-1750
C≡N	nitrile	2200-2250
C-H	alkane	2850-3100
N-H	amine, amide	3300-3500
O-H	carboxyl hydroxy	2500-3000 3200-3650





**Important values, constants and standards**

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ( $4.18 \text{ J g}^{-1} \text{ K}^{-1}$ )

The Periodic Table of Elements

		Group															
1	2											13	14	15	16	17	18
		<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">1 H hydrogen 1.0</div> <div style="border: 1px solid black; padding: 2px;">2 He helium 4.0</div> </div>															
		<div style="border: 1px solid black; padding: 5px;"> <b>Key</b>            atomic number            atomic symbol            name            relative atomic mass         </div>															
3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18		
Li lithium 6.9	Be beryllium 9.0	B boron 10.8	C carbon 12.0	N nitrogen 14.0	O oxygen 16.0	F fluorine 19.0	Ne neon 20.2	Na sodium 23.0	Mg magnesium 24.3	Al aluminium 27.0	Si silicon 28.1	P phosphorus 31.0	S sulfur 32.1	Cl chlorine 35.5	Ar argon 39.9		
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K potassium 39.1	Ca calcium 40.1	Sc scandium 45.0	Ti titanium 47.9	V vanadium 50.9	Cr chromium 52.0	Mn manganese 54.9	Fe iron 55.8	Co cobalt 58.9	Ni nickel 58.7	Cu copper 63.5	Zn zinc 65.4	Ga gallium 69.7	Ge germanium 72.6	As arsenic 74.9	Se selenium 79.0	Br bromine 79.9	Kr krypton 83.8
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb rubidium 85.5	Sr strontium 87.6	Y yttrium 88.9	Zr zirconium 91.2	Nb niobium 92.9	Mo molybdenum 95.9	Tc technetium —	Ru ruthenium 101.1	Rh rhodium 102.9	Pd palladium 106.4	Ag silver 107.9	Cd cadmium 112.4	In indium 114.8	Sn tin 118.7	Sb antimony 121.8	Te tellurium 127.6	I iodine 126.9	Xe xenon 131.3
55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs caesium 132.9	Ba barium 137.3	lanthanoids	Hf hafnium 178.5	Ta tantalum 180.9	W tungsten 183.8	Re rhenium 186.2	Os osmium 190.2	Ir iridium 192.2	Pt platinum 195.1	Au gold 197.0	Hg mercury 200.6	Tl thallium 204.4	Pb lead 207.2	Bi bismuth 208.0	Po polonium —	At astatine —	Rn radon —
87	88	89–103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118
Fr francium —	Ra radium —	actinoids	Rf rutherfordium —	Db dubnium —	Sg seaborgium —	Bh bohrium —	Hs hassium —	Mt meitnerium —	Ds darmstadtium —	Rg roentgenium —	Cn copernicium —	Nh nihonium —	Fl flerovium —	Mc moscovium —	Lv livermorium —	Ts tennessine —	Og oganeson —
57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74
La lanthanum 138.9	Ce cerium 140.1	Pr praseodymium 140.9	Nd neodymium 144.2	Pm promethium —	Sm samarium 150.4	Eu europium 152.0	Gd gadolinium 157.3	Tb terbium 158.9	Dy dysprosium 162.5	Ho holmium 164.9	Er erbium 167.3	Tm thulium 168.9	Yb ytterbium 173.1	Lu lutetium 175.0	102	103	104
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106
Ac actinium —	Th thorium 232.0	Pa protactinium 231.0	U uranium 238.0	Np neptunium —	Pu plutonium —	Am americium —	Cm curium —	Bk berkelium —	Cf californium —	Es einsteinium —	Fm fermium —	Md mendelevium —	No nobelium —	Lr lawrencium —	107	108	109

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.